


THIS CHILD HAS THE FOLLOWING ALLERGIES:

Name:

DOB:



Photo

Emergency contact details:

1)



2)



Child's Weight: Kg

Mild-moderate allergic reaction:

- Swollen lips, face or eyes
- Itchy / tingling mouth
- Hives or itchy skin rash
- Abdominal pain or vomiting
- Sudden change in behaviour

ACTION:

- Stay with the child, call for help if necessary
- Give antihistamine:
- Contact parent/carer (if vomited, can repeat dose)



Watch for signs of ANAPHYLAXIS (life-threatening allergic reaction):

- AIRWAY:** Persistent cough, hoarse voice, difficulty swallowing, swollen tongue
- BREATHING:** Difficult or noisy breathing, wheeze or persistent cough
- CONSCIOUSNESS:** Persistent dizziness / pale or floppy suddenly sleepy, collapse, unconscious

If ANY ONE of these signs are present:

1. Lie child flat. If breathing is difficult, allow to sit
2. Give EpiPen® or EpiPen® Junior
3. Dial 999 for an ambulance* and say ANAPHYLAXIS ("ANA-FIL-AX-IS")


If in doubt, give EpiPen®

After giving EpiPen:

1. Stay with child, contact parent/carer
2. Commence CPR if there are no signs of life
3. If no improvement **after 5 minutes**, give a further EpiPen® or alternative adrenaline autoinjector device, if available

*You can dial 999 from any phone, even if there is no credit left on a mobile. Medical observation in hospital is recommended after anaphylaxis.


How to give EpiPen®




Form fist around EpiPen® and PULL OFF BLUE SAFETY CAP



SWING AND PUSH ORANGE TIP against outer thigh (with or without clothing) until a click is heard



HOLD FIRMLY in place for 10 seconds



REMOVE EpiPen®. Massage injection site for 10 seconds

Keep your EpiPen device(s) at room temperature, do not refrigerate.

For more information and to register for a free reminder alert service, go to www.epipen.co.uk

Produced in conjunction with:



Additional instructions:

This is a medical document that can only be completed by the patient's treating health professional and cannot be altered without their permission.

This plan has been prepared by: _____

Hospital/Clinic: _____



Date: _____